



King's Research Portal

DOI:

[10.1002/ecs2.2625](https://doi.org/10.1002/ecs2.2625)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Kimmel, K., Dee, L., Tilman, D., Aubin, I., Boenisch, G., Catford, J. A., Kattge, J., & Isbell, F. (2019). Chronic fertilization and irrigation gradually and increasingly restructure grassland communities. *Ecosphere*, 10(3), [e02625]. <https://doi.org/10.1002/ecs2.2625>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Title: Chronic fertilization and irrigation gradually and increasingly restructure grassland communities

Authors: Kaitlin Kimmel^{1†}, Laura Dee², Dave Tilman^{1,3}, Isabelle Aubin⁴, Gerhard Boenisch⁵, Jane A. Catford^{6,7}, Jens Kattge^{5,8}, Forest Isbell¹

Author affiliations:

¹Department of Ecology, Evolution, and Behavior, University of Minnesota, St Paul, Minnesota 55108, USA

²Department of Fisheries, Wildlife and Conservation Biology, University of Minnesota, St Paul, Minnesota 55108, USA

³Bren School of Environmental Science & Management, University of California, Santa Barbara, California 93106, USA

⁴Great Lakes Forestry Centre, Canadian Forest Service, Natural Resources Canada, Sault Ste Marie, Ontario, P6A 2E5, Canada

⁵Max Planck Institute for Biogeochemistry, Hans Knöll Str. 10, 07745 Jena, Germany

⁶Department of Geography, King's College London, Strand, London WC2R 2LS, UK

⁷School of BioSciences, University of Melbourne, Vic 3010, Australia

⁸German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

†: Corresponding author: kimme099@umn.edu

Abstract

Scientists have known for over a century that resource addition can lead to species loss from plant communities. Recent studies have also shown that resource addition can substantially restructure communities by altering their functional and taxonomic composition – even when species richness remains unchanged. Understanding which aspects of community structure are impacted by different resources and over which timescales will provide insight for management decisions and may also elucidate which measures can act as early warning indicators for subsequent changes in the community. Here, we take advantage of a long-term factorial experiment to understand how grassland plant communities respond to a decade of nitrogen fertilization ($14 \text{ g N m}^{-2} \text{ year}^{-1}$) and irrigation ($25 \text{ mm water week}^{-1}$ during the growing season). After 10 years, fertilization and irrigation decreased species richness by 27% and 11%, while functional trait diversity decreased by 55% and 35%. Abundance-weighted functional distance between treatments and controls increased by 55% and 24%, respectively. We expected that abundance-weighted measures would shift before presence-absence based measures, but found limited evidence for this. Instead, our results suggest that species gains, which can occur quickly because they require the addition of only one individual, may serve as early indicators for subsequent community restructuring. Overall, both chronic nitrogen fertilization and irrigation tended to have gradual and increasing impacts on community structure, but the magnitude of these effects varied greatly depending on the aspect of community structure investigated. Further study will be needed to determine the extent to which our results can be generalized to other resources or sites in order to develop management strategies to maintain both taxonomic and functional trait diversity in the face of chronic resource changes.

Key words: Community structure, diversity, functional traits, nutrient addition, temporal trends, water addition

Introduction

Ecosystems are continually confronted with many long-term and directional changes in the supply of the major resources required by plants, including various nutrients (Galloway et al. 2004, IPCC 2014), water (Huntington 2006, Zhang et al. 2007), and CO₂ (IPCC 2014). Increased resource supply can restructure terrestrial plant communities (Lawes et al. 1882, Suding et al. 2005, Harpole et al. 2016) by altering functional and taxonomic richness, abundance, dominance, and composition. This community restructuring often occurs at the expense of species that would otherwise be dominant under low resource conditions, where they are able to drive resources down to levels at which their competitors can no longer replace themselves (Tilman 1982, Dybzinski and Tilman 2007). When resources are added, these low resource levels will be exceeded and another resource will instead become limiting, leading to shifts in species dominance and composition over time (Dybzinski and Tilman 2007, Isbell et al. 2013b). This raises several so far unanswered questions: (1) will changes in resource supply quickly restructure communities and have consistent and persistent effects, or instead restructure communities gradually and increasingly over time (Hillebrand et al. 2008, Smith et al. 2009, Mouillot et al. 2013b)? and (2) do some resources, or combinations of resources, have a greater influence on community structure than others (DeMalach et al. 2017)?

Community restructuring by chronic resource addition could have several possible temporal trends. Building on the framework proposed by Smith and colleagues (2015), we outline several distinct hypotheses about these temporal trends and their implications for communities (Fig. 1).

First, if the structure of a community is largely determined by factors other than resource supply (e.g., herbivory, pathogens, temperature; Cleland et al. 2011), community responses to increases in resource supply could vary from one year to the next, appearing temporally stochastic (H1–stochastic effect). On the other hand, if community structure is largely determined by resource limitation, community responses to increases in resource supply could grow, either linearly or non-linearly, over time as favored species increase in abundance and those disadvantaged decline (H2 – gradual effect). This would lead to increasing differences between regions which either experience this pressure or not. Alternatively, a community could restructure rapidly once the resource addition exceeds a threshold (H3- abrupt change; Bestelmeyer et al. 2011, Isbell et al. 2013). After reaching this threshold, the community may either continue to change or persist in its altered state. Finally, chronic resource addition may have a transient effect, where the community structure changes for a brief time before returning to its initial conditions (H4 – transient effect). These transient effects dynamics may happen if species have time to acclimate to the increased resource supply so that even if new species can become dominant over short timescales, original composition will recover over longer time scales (Reich et al. 2018).

Different aspects of community structure may respond differently to a change in resource supply or follow different temporal trends (Smith et al. 2009, Mouillot et al. 2013b). For example, metrics weighted by species' abundances, including many trait-based metrics, may be more responsive than presence-absence metrics, such as species richness (Smith et al. 2009, Mouillot et al. 2013b). However, many studies have only separately investigated either the effects of chronic nutrient addition on the species richness of grassland plant communities (Willems et al. 1993, Stevens et al. 2004, Clark and Tilman 2008, Harpole et al. 2016), or the divergence or convergence of species (Inouye and Tilman 1988, 1995, Houseman et al. 2008,

Avolio et al. 2015) and functional (Weiher et al. 1998, Laliberte et al. 2013, Mason et al. 2013, La Pierre and Smith 2015) composition. Using both taxonomic and functional trait metrics may not only provide insight into the temporal trends of community structure response but also give a more comprehensive picture of the extent of community restructuring.

Considering how both functional and taxonomic metrics respond through time can give insight into exactly what is driving changes perhaps concurrent changes in other metrics. For example, differences in species richness indicate changes in the number of species, but do not provide information regarding whether the species gained or lost are similar to or different from the original community. In contrast, changes in functional richness indicates that the species gained or lost were functionally unique (i.e., had different traits than the original species) or were replaced by functionally dissimilar or similar species (Cornwell et al. 2006, Villéger et al. 2008, Boersma et al. 2016). Yet, such patterns in functional richness could be generated both by concurrent changes in species richness or by species composition turnover (Fig. 2; e.g. Mayfield et al. 2010). Likewise, turnover in functional or taxonomic composition as well as shifts in relative abundances could occur without affecting functional or species richness indices (Fig. 2). Therefore, by using both functional trait metrics and traditional taxonomic community metrics we can more fully characterize how communities respond to different resource additions (Mason et al. 2005, Boersma et al. 2016) and the temporal trends of these responses (Smith et al. 2009, Mouillot et al. 2013b).

The objective of this study is to determine the extent to which temporal trends in community restructuring depend on the type of resource added and the metric of community structure investigated. Here, we examine temporal trends in the effects of chronic addition of water and nitrogen—two commonly limiting plant resources—on grassland plant community structure,

including multiple aspects of taxonomic and functional composition. We use a 10-year, full factorial water and nitrogen addition experiment at Cedar Creek (East Bethel, MN, USA)(Yang et al. 2018). We consider changes in several aspects of community structure: species richness and evenness, functional trait means and variance (functional dispersion, functional richness, and functional distance), community-weighted means of traits, and species and functional group abundances.

Methods

Study system

This study was conducted at Cedar Creek Ecosystem Science Reserve, East Bethel, MN, USA. Soils at Cedar Creek are characterized as nutrient-poor entisols derived from a glacial outwash sand plain (Tilman 1987). Average wet N deposition rates are $\sim 0.6 \text{ g N m}^{-2} \text{ year}^{-1}$ (58% NH_4 , 42% NO_3) (Clark and Tilman 2008). According to Köppen and Geiger classification (2006) the climate is characterized as cold continental with hot summer, but without dry season (Peel et al. 2007). The mean growing season (May – August) precipitation is approximately 420 mm, mean minimum growing season temperature is 12°C , and mean maximum growing season temperature is 25°C (1982-2016 period; <http://www.cedarcreek.umn.edu/research/data>). Plots are burned every year and fenced to prevent deer herbivory.

Experimental design

Thirty-six 9 x 9 m plots were planted with the same mixture of 32 grassland species in early spring of 1994 and again in 1995 (Tilman et al. 1997). Starting in 2007, plots were randomly assigned to one of six treatments, which included all combinations of two water treatments

(ambient and +25 mm/week during the growing season) and three nitrogen fertilization treatments (ambient, + 7 g N m⁻² year⁻¹), or +14 g N m⁻² year⁻¹). This water addition increases the total amount of water by ~80% during the average growing season, while the N fertilization rates are at levels relevant for understanding the impact of global wet deposition (Clark and Tilman 2008) and agricultural fertilization rates, respectively. Water was added to plots weekly from May to September via a sprinkler system, and nitrogen was added once in the beginning of each growing season as NH₄NO₃ (Farrior et al. 2013, Yang et al. 2018). Here, we show results from four of these treatments to simplify presentation (ambient, + water, + 14g N, and +water +14g N, n = 24), but similar results were obtained using both N addition treatments or grouping the N addition treatments together (Table S1). During 2007, 2008, 2010, 2015, and 2016, biomass from 4 strips, each 10 cm x 600 cm, was harvested, sorted to species, dried, and weighed. Non-planted species (weeds) were not removed from these plots, resulting in a total of 37 unplanted species observed across all plots and years. Overall, 96% of the biomass sampled across all plants and years was from the original 32 planted species.

Taxonomic metrics

We determined species richness (i.e. the number of species in a plot), inverse Simpson's evenness (Simpson 1949), and relative abundances of each species based on the harvested biomass from each plot. The biomass of the ten most abundant species over all plots and experiment years comprised ~82% of all biomass sampled. These ten species were: *Andropogon gerardii* (29.4%), *Sorghastrum nutans* (9.8%), *Schizachyrium scoparium* (8.5%), *Poa pratensis* (8.4%), *Lupinus perennis* (6.8%), *Dalea purpurea* (5.6%), *Coreopsis palmata* (3.9%), *Bouteloua gracilis* (3.1%), *Liatiris aspera* (3%), *Solidago rigida* (2.9%). For analyses of individual species'

biomass, the square-root of the biomass was used as the response variable in order to meet model assumptions of normality.

Functional traits

We characterized each species by four different functional traits which can be indicative of success when resource supplies increase (Craine et al. 2002, Sandel et al. 2010, La Pierre and Smith 2015). We included plant height (m) because it is associated with the ability to compete for light resources (Gaudet and Keddy 1988, Westoby 1998) and two leaf economics traits (specific leaf area (SLA), $\text{mm}^2 \text{mg}^{-1}$ and leaf tissue nitrogen per dry mass (leaf N), mg g^{-1}) because they are associated with the trade-off between acquisitive, fast-growing, but short-lived leaves and conservative, slow-growing, but long-lived leaves (Wright et al. 2004, Reich 2014). We also included photosynthetic pathway (C3/C4) because it is related to seasonality and resource use efficiency. Trait data were obtained from Cedar Creek datasets (<http://www.cedarcreek.umn.edu/research/data>) and the global plant trait database – TRY (Kattge et al. 2011; www.try-db.org, see Table S2 for a summary of data used and associated references). We averaged each trait over all records for each species when more than one observation existed in the databases. Over half of the species had more than 10 observations, and dominant species had from 50 – 650 observations. Species missing information for more than two traits were excluded from trait-based analyses, though we were able to characterize between 87 – 100% of species and 89 - 100% of biomass present in each treatment and year with available data (Table S3). While species may shift their trait values in response to changing environmental conditions (Lepš et al. 2011), species reordering, gains and losses are expected to dominate community responses over the longer timescales of our experiment (Smith et al. 2009).

Thus, characterizing species by average values captures changes in the *types* of species but not intraspecific trait responses.

Functional trait metrics

We calculated three different functional trait metrics; each testing a specific aspect of functional composition response to increases in resource supply. First, functional richness measures the amount of trait space occupied by a community based on the presence or absence of traits, regardless of their relative abundance in the community (Mason et al. 2005, Cornwell et al. 2006). Significant decreases in functional richness indicate convergence toward a similar trait composition (Laliberte et al. 2013, Mason et al. 2013, Boersma et al. 2016). Functional richness is calculated as the convex hull of the multivariate trait space, the multidimensional space where each axis corresponds to a trait. Second, functional dispersion quantifies the spread of traits in the community based on both presence and relative abundances (Laliberte and Legendre 2010). Significant changes in functional dispersion, without parallel changes in functional richness, can indicate shifts in trait dominance and evenness (Boersma et al. 2016). Functional dispersion is calculated as the abundance-weighted average of the distance that each species (trait combination) is away from an abundance-weighted centroid of the community (i.e., within a plot) (Laliberte and Legendre 2010). Third, functional distance indicates whether treated plots are located in different areas of trait space compared to ambient plots (Boersma et al. 2016). Functional distance is calculated as the distance between the centroids of each pair of plots to determine whether distances between pairs of plots within the same treatments are less than those between treatments (*see* Boersma et al. 2016). In our analyses, increases in functional distance through time indicate shifts in species abundance and species turnover, such that treatment plots

are moving further away from ambient plots in trait space. For each trait, a community-weighted mean and a non-weighted mean was calculated for each plot at each year to investigate which *types* of species (e.g., do we tend to move towards species with high tissue nutrient concentration?) were responding based on abundance and presence/absence. Lastly, we also grouped species into four functional groups based on broadly similar physiology and morphology - C4 grasses, C3 grasses, legumes, and all other non-legume forbs- to classify how these groups respond to our treatments. These functional groups have been previously shown to be responsive to long-term nutrient addition (Tilman et al. 1997, Reich et al. 2004).

We calculated functional richness and dispersion for each plot within each year using the FD package in R (Laliberte and Legendre 2010, Laliberté et al. 2015). We did the analyses with both z -transformed and untransformed trait values. There were no qualitative differences between the two methods, so we present data using untransformed trait values. We set $m = 4$ to compute functional richness since we characterized each species by 4 different traits. We calculated functional distance based on abundance-weighted and non-abundance-weighted distances between plots to assess the extent to which results depended on shifts in relative abundance, rather than changes in the presence or absence of species (Boersma et al. 2016); see *Statistical analyses* for a discussion on how this was modeled.

Statistical analyses

To determine how functional and taxonomic metrics responded to treatments, our estimated models considered (1) the three-way interaction of each treatment and experiment duration, (2) all pair-wise interactions, (3) the main effect of each treatment and study duration. The models also accounted for effects of average annual biomass production and the standardized

precipitation-evapotranspiration index (SPEI, Fig. S1). We build off of previous frameworks for testing temporal trends (Smith et al. 2009, 2015), but determine temporal trends from statistical models which represent our hypothesized relationships and AIC model selection. Reflecting our hypotheses about temporal trends, we consider three candidate functional forms for the interaction between nutrient treatment, irrigation treatment, and experimental duration: linear (gradual response; H2), log-linear (gradual response; H2), and quadratic (transient response; H4), along with year as a factor (stochastic response; H1). If there were only main effects for the treatments, we considered this a threshold response (H3). We included the average biomass production each year and the standardized precipitation-evapotranspiration index (SPEI; Vicente-Serrano et al. 2010) for the 5-month growing season (April – August) to control for their effects and get more precise estimates of the treatments effects on measures of community structure. Model selection was based on the lowest AIC for the results shown, with $\Delta AIC > 2$ (Table S1). If the 3-way interaction between the two treatments and year was not significant, it was removed for parsimony (Crawley 2013). We removed SPEI and biomass only if this improved AIC values by $\Delta AIC > 2$, which occurred only for SLA. Reflecting our hypotheses about temporal trends, if the quadratic form fit best, but had no treatment by year² interactions (e.g. no transient effect), the next best model was used.

To account for plot-by-plot variability, we considered models that included random slopes for each treatment and random intercepts for each plot. After model selection, we used random intercept only models, which had the lowest AIC values ($\Delta AIC > 5$). To account for the repeated measures through time, we used compound symmetry covariance structures (Pinheiro and Bates 2000). All models selected had the compound symmetry structure. We were unable to consider other temporal covariance structures, such as a first-order autoregressive structure,

because data were not collected every year. All analyses were completed in R v 3.3.3 (R Core Team 2017) using the lme function in the nlme package (Pinheiro et al. 2017).

We modeled both abundance-weighted and non-abundance-weighted functional distance as outlined in Boersma et al (2016) using the MCMCglmm package in R (Hadfield 2010). We compared ambient plots to all treatment plots to determine whether increased resource supply causes communities to move to a different area of trait space. We constructed a trait matrix that included all species present for all years and an abundance matrix for all years to calculate centroids and distances between plots in multidimensional trait space (see Boersma et al 2016). We then selected for only within year distances. Comparisons were made between ambient plots and all treatment plots to determine how treatments impacted trait shifts. To account for the multiple comparisons between plots within each year, we modeled random effects as a multi-membership model where variance is equally distributed across all pairs of plots within each year. We also included average biomass for each year and the 5-month SPEI in our models to more accurately predict the impacts of the specific treatments on functional distance. The best model for the abundance-weighted distances included SPEI and biomass (Fig. S1). The best model for the natural logarithm of non-abundance-weighted distances included year as a factor (Fig. S2).

Results

Overall, we found that fertilization and irrigation caused most measures of community structure to gradually deviate from ambient plots, and these effects increased through time (Table 1, H2). Over the duration of the experiment, fertilization and irrigation impacted a similar number of community metrics. Under fertilization treatments 11 measures of community structure

responded with continuous gradual changes over time, four displayed stochastic (but directional) changes, four changed abruptly, and one showed transient change. Under irrigation treatments, 12 measures of community structure exhibited continuous changes, five changed abruptly, three showed transient change, and one responded stochastically (Table 1). The trends in the community structure measures were often mirrored by species-specific and individual functional trait responses, although not always.

Taxonomic restructuring

Fertilization rapidly increased species richness in the first year, but had a continuous negative impact on species richness and Simpson's diversity over the duration of this experiment (Fig. 3A-B, Fig. S1; H2). Overall, fertilization decreased species richness by 27% and Simpson's diversity by 45%. The effect of irrigation on species richness also increased over the duration of the experiment, amounting to an 11% decrease (Fig. 3B, Fig. S1; H2). In contrast, irrigation did not have a significant effect on Simpson's diversity (Fig. 3A, Fig. S1). Fertilization marginally decreased evenness by 15% within the first year, and this effect persisted at approximately the same magnitude during subsequent years (Fig. 3C, Fig. S1; H3).

Functional restructuring

Fertilization caused a stochastic decrease in functional richness over the duration of the experiment; there was an overall decline in functional richness resulting in a 55% decrease compared to ambient plots, but the magnitude of the effect was variable from year to year (Fig. 3D, Fig. S1; H1). Fertilization also caused functional composition to become increasingly different from ambient plots over time when considering abundance weighted distances (Fig. 3F,

Fig. S1; H2) and caused a stochastic divergence when considering non-abundance-weighted distances (Fig. S2; H1). Thus, fertilization caused both a contraction of trait space and shifted the region of trait space occupied by each community (Fig. 2D). Irrigation also caused plots to diverge from ambient plots through time, but only when considering abundance-weighted distances (Fig. 3F, Fig. S1). Functional dispersion did not respond to any of the treatments and remained relatively consistent through time (Fig. 3E, Fig. S1).

Community-level functional responses

Functional group composition was influenced by both nitrogen addition and irrigation (Fig. 4 and 5, Table S4). Fertilization had a rapid and persistent effect on the proportion of the community represented by legumes (negative) and C3 grasses (positive; H3), and had gradual continuous effects on the proportion of non-legume forbs (negative) and C4 grasses (positive; Fig. 4A-D; H1). The decrease in non-legume forbs was mirrored by a continuous decrease in the proportion of the community weighted mean proportion of C3 species and height (Fig. 4G-H, Table S6). Irrigation had positive, rapid and persistent impacts on the proportion of C4 grasses in the community (H3), decreased the proportion non-legume forbs and C3 grasses through time (H1), and increased the proportion of legumes through time (Fig. 4A-D, H1). Plots that were both fertilized and irrigated decreased in non-legume forbs less than expected from the additive effects of each treatment separately (Fig 4C). Changes in functional groups were mirrored by rapid decreases in community weighted means of leaf N and proportion of C3 species (Fig. 4F & H, Table S6). Most of the functional groups and community-weighted traits had rapid responses to at least one of the treatments in the first year of the experiment (Fig. 4, Table 1).

Species-specific responses

Irrigation caused changes in the biomass of eight of the ten most dominant species whereas nitrogen caused changes in biomass of seven of these ten species (Fig.5, Fig. S3). The most common response was a linear change through time (six of ten, H2). Species-specific responses reflected the trait and functional group responses. For example, fertilization had an increasingly positive effect on *Andropogon gerardii* biomass, a C4 species with low tissue N and moderately high SLA. In contrast, fertilization had an increasingly negative effect on *Lupinus perennis* biomass, a C3 legume with high tissue nitrogen (Fig. 5; see Fig. S3 for more details).

Discussion

Here, we show that temporal trends in the plant community response to chronic resource addition depend on both the resource and the specific aspect of community structure under consideration. In this experiment, we find that nitrogen fertilization and irrigation gradually and increasingly impacted many aspects of community structure. Few metrics had stochastic or transient responses, while some had abrupt responses. Increasing species loss over time with nutrient addition is consistent with other previous studies both at our study site (Clark and Tilman 2008, Isbell et al. 2013a) and globally (Stevens et al. 2004, Maskell et al. 2010, Harpole et al. 2016). We further show that several aspects of community structure had gradual and increasing responses to fertilization over time. Globally, the short-term effects of water addition on species richness is known to vary considerably across study sites, but has a positive effect on average on species richness (DeMalach et al. 2017). Our results suggest that the long-term effects of increased water availability on community structure may be discernable from short-term studies because gradual, continuous responses were common. While many metrics had this

gradual and continuous response to chronic resource addition, not all had the same temporal trends.

Using multiple metrics of plant community composition allowed for a more comprehensive understanding of community restructuring and allowed us to tease apart changes in certain aggregate metrics. We found that both species richness and functional richness were declining in fertilized treatments, providing evidence that fertilization causes not only species loss, but also loss of functionally unique species. This effect is consistent with results from other sites (Suding et al. 2005) and provides evidence that the species lost as a result of resource addition were those with distinctive trait combinations. The loss of distinctive trait combinations can reduce ecosystem stability and affect functioning, potentially reducing the range of conditions in which the community can maintain high levels of productivity (Isbell et al. 2011, Mouillot et al. 2013a, Violle et al. 2017). Alternatively, their disappearance could increase ecosystem stability if highly stable species persist (Polley et al. 2007). Further, fertilized and irrigated plots occupy a different region of trait space compared to ambient plots. Specifically, we found that both fertilization and irrigation increased the abundance of species with higher SLA and decreased the abundance of C3 species, whereas irrigation shifted communities towards species with lower leaf nitrogen, which were consistent with some other studies (Sandel et al. 2010, La Pierre and Smith 2015, but see Isbell et al. 2013b). This contraction of trait space may indirectly impact certain ecosystem functions like productivity through time (Isbell et al. 2013a), if, for example, it reduces complementarity between species (Loreau and Hector 2001, Fargione et al. 2007). It could also potentially increase productivity by selecting for more productive species in the new environmental context (Roscher et al. 2012, Cadotte 2017). More studies are needed to understand the consequences of these trait shifts on ecosystem functioning.

We expected that abundance-weighted or functional trait metrics would shift before presence-absence based measures (Hillebrand et al. 2008, Mouillot et al. 2013b), but found limited evidence for this. Instead, our results also suggest that species gains, which can occur quickly because they require the addition of only one individual, may serve as early indicators for subsequent species losses, which often take more time because they require the loss of the last individual. While we found that many abundance-weighted metrics, including Simpson's diversity, evenness, and community-weighted means, can have rapid responses to chronic resource addition these metrics did not necessarily deviate much from ambient temporal trends through time, indicating limits to their sensitivity. Conversely, we saw a rapid increase in both species richness and functional richness within the first year that indicates functionally unique species were added to the community. Rather than turnover, these species were lost over the duration of experiment because many of the dominant species and functional groups tended to increase over time. Further, many of the functional groups responded rapidly to irrigation, and community weighted means of traits responded rapidly to both fertilization and irrigation. This suggests that the early-indicator metrics may differ from resource to resource. Nevertheless, it does not seem that abundance-weighted or functional trait metrics captured changes in community composition sooner than taxonomic measures.

Nitrogen and water are expected to restructure the community in different ways because of how these treatments were applied and how these resources interact within the system. In this experiment, are more comparable to agricultural levels of nitrogen fertilization than rates of atmospheric N deposition. Irrigation treatment levels increased water by 80% compared to average ambient precipitation levels over the past 30 years. Further, nitrogen was applied once at the beginning of the growing season whereas irrigation occurred weekly throughout the growing

season, so differences in resource addition frequency may cause differences in how plants compete for these resources. Irrigation likely reduced water stress throughout the season whereas nitrogen addition may have had a disproportionate impact on species establishment early in the growing season. Furthermore, nitrogen can accumulate in the system over multiple years (Isbell et al. 2013b) whereas although irrigation can alleviate water stress throughout the period of application, it is not retained by the sandy soils at our site, limiting the cumulative effect of chronic application. The resulting difference in effective resource availability over time, may help to explain why the effects of nitrogen were greater in magnitude at our site. Cedar Creek is strongly nitrogen limited, and adding nitrogen consistently increases aboveground plant biomass, reduces species richness, and shifts dominance (Tilman 1982, Clark and Tilman 2008, Isbell et al. 2013a). Nitrogen is therefore expected to have a greater effect on community structure than water availability, which is not a limitation at our site, except perhaps during droughts, rainfall being higher compared to many other grassland sites (Cleland et al. 2013). It follows that more xeric grasslands may have greater responses to water addition (Cleland et al. 2013, Wilcox et al. 2017). These experimental treatments could impact sites with different climates differently than what we found here, but we expect that gradual, continuous responses should dominate response to chronic pressures.

Environmental changes cause shifts in the functional structure of a community as the result of species replacement and within-species variation lead (Lepš et al. 2011). In this paper, we focused on the trait variability between species, i.e. using one average trait values for each species for all plots samples, variation between species is thought to be greater than variation within a single species (*but see* Albert et al. 2010, Moran et al. 2016). Further, we were interested in long term trends which are primarily related to species replacement (i.e. changes in

species composition or abundance). However, the use of average trait values can mask within species variation, i.e., the variation between individuals exposed to different environmental conditions (Smith et al. 2009). Therefore, using average trait values may conceal some changes in community structure that could otherwise be detected by measuring traits for every species in every treatment during every year (Jung et al. 2010, Violle et al. 2012). However, these intraspecific differences are expected to be more important in the case of short term assessment (Smith et al. 2009, Auger and Shipley 2013) when the taxonomic structure of the community is more similar to previous states. There is still clearly a need to sort out how important intraspecific trait variation is in determining community structure and over what time scales it is relevant.

Overall, we found that chronic resource additions resulted in continuous, gradual changes in communities, consistent with H2, and that nitrogen and irrigation impacted a similar number of metrics. For metrics that responded to both fertilization and irrigation, fertilization had much a larger effect size. While gradual and increasing changes were the most common, we found evidence for each of our hypothesized trends under both treatments regimes. Because of this array of responses, further research is needed to determine the extent to which our results can be generalized to other resources or sites. Understanding these temporal trends at different sites and for different resources will help inform our expectations for when and how much chronic resource addition will modify communities. We will also be able to show what metrics are early indicators of future change. Thus, discerning exactly when and where we expect these temporal trends to occur will help us develop management strategies to maintain both taxonomic and functional trait diversity in the face of these chronic resource changes.

438 **Author Contributions:**

439 KK – Conceived the project and performed the data analysis and led the writing
440 LD – Contributed to conceptualization of temporal trends and hypotheses, conceptual figures,
441 analyses, and writing
442 DT – Designed experiment, contributed to writing
443 IA - Contributed to plant trait data and writing
444 GB - Contributed to plant trait data and writing
445 JC – Contributed to plant trait data and writing
446 JK – Contributed to plant trait data and writing
447 FI – Contributed to analyses and writing

448

449 **Acknowledgements:**

450 We thank Troy Mielke, Dan Bahaiddin, Kally Worm, and many summer interns for their
451 assistance with this research. This study has been supported by the TRY initiative on plant traits
452 (<http://www.try-db.org>). The TRY initiative and database is hosted, developed and maintained
453 by J. Kattge and Bonisch (Max Planck Institute for Biogeochemistry, Jena, Germany). TRY is
454 currently supported by DIVERSITAS/Future Earth and the German Centre for Integrative
455 Biodiversity Research (iDiv) Halle-Jena-Leipzig. This work was supported by grants from the
456 US National Science Foundation Long-Term Ecological Research Program (LTER) including
457 DEB-0620652 and DEB-1234162. Further support was provided by the Cedar Creek Ecosystem
458 Science Reserve and the University of Minnesota.

459

460

Literature cited

- Albert, C. H. C. H., W. Thuiller, N. G. Yoccoz, A. Soudant, F. Boucher, P. Saccone, and S. Lavorel. 2010. Intraspecific functional variability: extent, structure and sources of variation. *Journal of Ecology* 98:604–613.
- Auger, S., and B. Shipley. 2013. Inter-specific and intra-specific trait variation along short environmental gradients in an old-growth temperate forest. *Journal of Vegetation Science* 24:419–428.
- Avolio, M. L., K. J. La Pierre, G. R. Houseman, S. E. Koerner, E. Grman, F. I. Isbell, D. S. Johnson, and K. R. Wilcox. 2015. A framework for quantifying the magnitude and variability of community responses to global change drivers. *Ecosphere* 6:1–14.
- Bestelmeyer, B. T., A. M. Ellison, W. R. Fraser, K. B. Gorman, S. J. Holbrook, C. M. Laney, M. D. Ohman, D. P. C. Peters, F. C. Pillsbury, A. Rassweiler, R. J. Schmitt, and S. Sharma. 2011. Analysis of abrupt transitions in ecological systems. *Ecosphere* 2:art129.
- Boersma, K. S., L. E. Dee, S. J. Miller, M. T. Bogan, D. A. Lytle, and A. I. Gitelman. 2016. Linking multidimensional functional diversity to quantitative methods: a graphical hypothesis-evaluation framework. *Ecology* 97:583–593.
- Cadotte, M. W. 2017. Functional traits explain ecosystem function through opposing mechanisms. *Ecology Letters*:1–8.
- Clark, C. M., and D. Tilman. 2008. Loss of plant species after chronic low-level nitrogen deposition to prairie grasslands. *Nature* 451:712–5.
- Cleland, E. E., C. M. Clark, S. L. Collins, J. E. Fargione, L. Gough, K. L. Gross, S. C. Pennings, and K. N. Suding. 2011. Patterns of trait convergence and divergence among native and exotic species in herbaceous plant communities are not modified by nitrogen enrichment.

484 Journal of Ecology 99:1327–1338.

485 Cleland, E. E., S. L. Collins, T. L. Dickson, E. C. Farrer, K. L. Gross, L. A. Gherardi, L. M.

486 Hallett, R. J. Hobbs, J. S. Hsu, L. Turnbull, and K. N. Suding. 2013. Sensitivity of grassland

487 plant community composition to spatial versus temporal variation in precipitation. *Ecology*

488 94:1687–1696.

489 Cornwell, W. K., D. W. Schwilk, and D. D. Ackerly. 2006. A trait-based test for habitat filtering:

490 Convex hull volume. *Ecology* 87:1465–1471.

491 Craine, J. M., D. Tilman, D. Wedin, P. B. Reich, and M. Tjoelker. 2002. Functional traits,

492 productivity and effects on nitrogen cycling of 33 grassland species. *Functional Ecology*

493 16:563–574.

494 Crawley, M. J. 2013. *The R Book*. Second edition. Wiley, Chichester, West Sussex, United

495 Kingdom.

496 DeMalach, N., E. Zaady, and R. Kadmon. 2017. Contrasting effects of water and nutrient

497 additions on grassland communities: A global meta-analysis. *Global Ecology and*

498 *Biogeography* 26:983–992.

499 Dybzinski, R., and D. Tilman. 2007. Resource use patterns predict long-term outcomes of plant

500 competition for nutrients and light. *The American Naturalist* 170:305–318.

501 Fargione, J., D. Tilman, R. Dybzinski, J. Hille Ris Lambers, C. Clark, W. S. Harpole, J. M. H.

502 Knops, P. B. Reich, and M. Loreau. 2007. From selection to complementarity: shifts in the

503 causes of biodiversity- productivity relationships in a long-term biodiversity experiment.

504 *Proceedings of the Royal Society B*.

505 Farrior, C. E., D. Tilman, R. Dybzinski, P. B. Reich, S. Levin, and S. Pacala. 2013. Resource

506 limitation in a competitive context determines complex plant responses to experimental

507 resource additions. *Ecology* 94:2505–2517.

508 Galloway, J. N., F. J. Dentener, D. G. Capone, E. W. Boyer, R. W. Howarth, S. P. Seitzinger, G.
509 P. Asner, C. C. Cleveland, P. A. Green, E. A. Holland, D. M. Karl, A. F. Michaels, J. H.
510 Porter, A. R. Townsend, and C. J. Vorosmarty. 2004. Nitrogen cycles: past, present and
511 future. *Biogeochemistry* 70:153–226.

512 Gaudet, C. L., and P. A. Keddy. 1988. A comparative approach to predicting competitive ability
513 from plant traits. *Nature* 334:242.

514 Hadfield, J. D. 2010. MCMC methods for multi-response generalized linear mixed models: the
515 MCMCglmm R package. *Journal of Statistical Software* 33:1–22.

516 Harpole, W. S., L. L. Sullivan, E. M. Lind, J. Firn, P. B. Adler, E. T. Borer, J. Chase, P. A. Fay,
517 Y. Hautier, H. Hillebrand, A. S. MacDougall, E. W. Seabloom, R. Williams, J. D. Bakker,
518 M. W. Cadotte, E. J. Chaneton, C. Chu, E. E. Cleland, C. D’Antonio, K. F. Davies, D. S.
519 Gruner, N. Hagenah, K. Kirkman, J. M. H. Knops, K. J. La Pierre, R. L. McCulley, J. L.
520 Moore, J. W. Morgan, S. M. Prober, A. C. Risch, M. Schuetz, C. J. Stevens, and P. D.
521 Wragg. 2016. Addition of multiple limiting resources reduces grassland diversity. *Nature*
522 537:1–9.

523 Hillebrand, H., D. Bennett, and M. W. Cadotte. 2008. Consequences of dominance: A review of
524 evenness effects on local and regional ecosystem processes. *Ecology* 89:1510–1520.

525 Houseman, G. R., G. G. Mittelbach, H. L. Reynolds, and K. L. Gross. 2008. Perturbations alter
526 community convergence, divergence, and formation of multiple community states. *Ecology*
527 89:2172–2180.

528 Huntington, T. G. 2006. Evidence for intensification of the global water cycle: Review and
529 synthesis. *Journal of Hydrology* 319:83–95.

530 Inouye, R. S., and D. Tilman. 1988. Convergence and divergence of old-field plant communities
 531 along experimental nitrogen gradients. *Ecology* 69:995–1004.

532 Inouye, R. S., and D. Tilman. 1995. Convergence and divergence of old-field vegetation after 11
 533 yr of nitrogen addition. *Ecology* 76:1872–1887.

534 IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and
 535 III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.

536 Isbell, F. I., V. Calcagno, A. Hector, J. Connolly, W. S. Harpole, P. B. Reich, M. Scherer-
 537 Lorenzen, B. Schmid, D. Tilman, J. van Ruijven, A. Weigelt, B. J. Wilsey, E. S. Zavaleta,
 538 and M. Loreau. 2011. High plant diversity is needed to maintain ecosystem services. *Nature*
 539 477:199–202.

540 Isbell, F. I., P. B. Reich, D. Tilman, S. E. Hobbie, S. Polasky, and S. Binder. 2013a. Nutrient
 541 enrichment, biodiversity loss, and consequent declines in ecosystem productivity. *PNAS*
 542 110:11911–6.

543 Isbell, F. I., D. Tilman, S. Polasky, S. Binder, and P. Hawthorne. 2013b. Low biodiversity state
 544 persists two decades after cessation of nutrient enrichment. *Ecology letters* 16:454–60.

545 Jung, V., C. Violle, C. Mondy, L. Hoffmann, and S. Muller. 2010. Intraspecific variability and
 546 trait-based community assembly. *Journal of Ecology* 98:1134–1140.

547 Kattge, J., S. Díaz, S. Lavorel, I. C. Prentice, P. Leadley, G. Bönsch, E. Garnier, M. Westoby, P.
 548 B. Reich, I. J. Wright, J. H. C. Cornelissen, M. Violle, C S. P. Harrison, P. M. v. Bodegom,
 549 T. Reichstein, B. J. Enquist, N. A. Soudzilovskaia, D. D. Ackerly, M. Anand, O. Atkin, M.
 550 Bahn, D. E. R. Baker, D. Baldocchi, R. Bekker, C. Blanco, B. Blonder, W. J. Bond, R.
 551 Bradstock, D. C. Bunker, F. Casanoves, J. Cavender-Bares, J. Q. Chambers, F. S. Chapin, J.
 552 Chave, M. E. W. K. Cornwell, J. M. Craine, B. H. Dobrin, L. Duarte, W. Durka, J. Elser, G.

553 Esser, D. F. W. F. Fagan, J. Fang, F. Fernández-Méndez, A. Fidelis, B. Finegan, O. Flores,
 554 H. Ford, S. G. T. Freschet, N. M. Fyllas, R. V. Gallagher, W. A. Green, A. G. Gutierrez, T.
 555 Hickler, M. Higgins, J. G. Hodgson, A. Jalili, S. Jansen, C. Joly, A. J. Kerkhoff, D. Kirkup,
 556 K. Kitajima, M. Kleyer, S. Klotz, J. M. H. Knops, K. Kramer, I. Kühn, H. Kurokawa, D.
 557 Laughlin, T. D. Lee, P. Leishman, F. Lens, T. Lenz, S. L. Lewis, J. Lloyd, J. Llusià, F.
 558 Louault, S. Ma, M. D. Mahecha, S. N. Manning, T. Massad, B. Medlyn, J. Messier, A. T.
 559 Moles, S. C. Müller, K. Nadrowski, J. Ü. Niinemets, S. Nöllert, A. Nüske, R. Ogaya, J.
 560 Oleksyn, V. G. Onipchenko, Y. Onoda, O. L. Ordoñez, G. Overbeck, W. A. Ozinga, S.
 561 Patiño, S. Paula, J. G. Pausas, J. Peñuelas, S. Phillips, V. Pillar, H. Poorter, L. Poorter, P.
 562 Poschlod, A. Prinzing, R. Proulx, A. Rammig, E. Reinsch, B. Reu, L. Sack, B. Salgado-
 563 Negret, J. Sardans, S. Shiodera, B. Shipley, A. Siefert, E. Sosinski, J.-F. Soussana, E.
 564 Swaine, N. Swenson, K. Thompson, P. Thornton, M. Waldram, and C. W. Weiher, M.
 565 White, S. White, S. J. Wright, B. Yguel, S. Zaehle, A. E. Zanne. 2011. TRY – a global
 566 database of plant traits. *Global Change Biology* 17:2905–2935.

567 Laliberte, E., and P. Legendre. 2010. A distance-based framework for measuring functional
 568 diversity from multiple traits. *Ecology* 91:299–305.

569 Laliberté, E., P. Legendre, and Bill Shipley. 2015. Measuring functional diversity (FD) from
 570 multiple traits, and other tools for functional ecology.

571 Laliberte, E., D. A. Norton, and D. Scott. 2013. Contrasting effects of productivity and
 572 disturbance on plant functional diversity at local and metacommunity scales. *Journal of*
 573 *Vegetation Science* 24:834–842.

574 Lawes, J. B., J. H. Gilbert, and M. T. Masters. 1882. Agricultural, botanical, and chemical results
 575 of experiments on the mixed herbage of permanent meadow, conducted for more than

576 twenty years in succession on the same land. Part II. The botanical results. Royal Society of
577 London Philosophical Transactions Series I 173:1181–1413.

578 Lepš, J., F. de Bello, P. Šmilauer, and J. Doležal. 2011. Community trait response to
579 environment: Disentangling species turnover vs intraspecific trait variability effects.
580 *Ecography* 34:856–863.

581 Loreau, M., and a Hector. 2001. Partitioning selection and complementarity in biodiversity
582 experiments. *Nature* 412:72–6.

583 Maskell, L. C., S. M. Smart, J. M. Bullock, K. Thompson, and C. J. Stevens. 2010. Nitrogen
584 deposition causes widespread loss of species richness in British habitats. *Global Change*
585 *Biology* 16:671–679.

586 Mason, N. W. H., F. De Bello, D. Mouillot, S. Pavoine, and S. Dray. 2013. A guide for using
587 functional diversity indices to reveal changes in assembly processes along ecological
588 gradients. *Journal of Vegetation Science* 24:794–806.

589 Mason, N. W. H., D. Mouillot, W. G. Lee, J. B. Wilson, and H. Setälä. 2005. Functional
590 richness, functional evenness and functional divergence: The primary components of
591 functional diversity. *Oikos* 111:112–118.

592 Mayfield, M. M., S. P. Bosner, J. W. Morgan, I. Aubin, S. McNamara, and P. A. Vesk. 2010.
593 What does species richness tell us about function trait diversity? Predictions and evidence
594 of species and function trait diversity to land-use change. *Global Ecology and*
595 *Biogeography* 19:423–431.

596 Moran, E. V, F. Hartig, and D. M. Bell. 2016. Intraspecific trait variation across scales:
597 implications for understanding global change responses. *Global Change Biology* 22:137–
598 150.

599 Mouillot, D., D. R. Bellwood, C. Baraloto, J. Chave, R. Galzin, M. Harmelin-Vivien, M.
 600 Kulbicki, S. Lavergne, S. Lavorel, N. Mouquet, C. E. T. Paine, J. Renaud, and W. Thuiller.
 601 2013a. Rare species support vulnerable functions in high-diversity ecosystems. *PLoS*
 602 *Biology* 11.
 603 Mouillot, D., N. A. J. Graham, S. Villegger, N. W. H. Mason, and D. R. Bellwood. 2013b. A
 604 functional approach reveals community responses to disturbances. *Trends in Ecology and*
 605 *Evolution* 28:167–177.
 606 Peel, M. C., B. L. Finlayson, and T. A. McMahon. 2007. Updated world map of the Köppen–
 607 Geiger climate classification. *Hydrology and Earth Systems Science* 11:1633–1644.
 608 La Pierre, K. J., and M. D. Smith. 2015. Functional trait expression of grassland species shift
 609 with short- and long-term nutrient additions. *Plant Ecology* 216:307–318.
 610 Pinheiro, J., D. Bates, S. DebRoy, D. Sarkar, and R Core Team. 2017. nlme: Linear and
 611 nonlinear mixed effects models.
 612 Pinheiro, J. C., and D. M. Bates. 2000. Correlation structures for modeling dependence. Pages
 613 226–248 *Mixed-effects models in S and S-Plus*. Springer, New York.
 614 Polley, H. W., B. J. Wilsey, and J. D. Derner. 2007. Dominant species constrain effects of
 615 species diversity on temporal variability in biomass production of tallgrass prairie. *Oikos*
 616 116:2044–2052.
 617 R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for
 618 Statistical Computin, Vienna, Austria.
 619 Reich, P. B. 2014. The world-wide “fast-slow” plant economics spectrum: A traits manifesto.
 620 *Journal of Ecology* 102:275–301.
 621 Reich, P. B., S. E. Hobbie, T. D. Lee, and M. A. Pastore. 2018. Unexpected reversal of C3 versus

622 C4 grass response to elevated CO₂ during a 20-year field experiment. *Science* 360:317–
623 320.

624 Reich, P. B., D. Tilman, S. Naeem, D. S. Ellsworth, J. Knops, J. Craine, D. Wedin, and J. Trost.
625 2004. Species and functional group diversity independently influence biomass accumulation
626 and its response to CO₂ and N. *PNAS* 101:10101–10106.

627 Roscher, C., J. Schumacher, M. Gubsch, A. Lipowsky, A. Weigelt, N. Buchmann, B. Schmid,
628 and E. D. Schulze. 2012. Using plant functional traits to explain diversity-productivity
629 relationships. *PLoS ONE* 7.

630 Sandel, B., L. J. Goldstein, N. J. B. Kraft, J. G. Okie, M. I. Shuldman, D. D. Ackerly, E. E.
631 Cleland, and K. N. Suding. 2010. Contrasting trait responses in plant communities to
632 experimental and geographic variation in precipitation. *New Phytologist* 188:565–575.

633 Simpson, E. H. 1949. Measurement of Diversity. *Nature* 163:688.

634 Smith, M. D., A. K. Knapp, and S. L. Collins. 2009. A framework for assessing ecosystem
635 dynamics in response to chronic resource alterations induced by global change. *Ecology*
636 90:3279–3289.

637 Smith, M. D., K. J. La Pierre, S. L. Collins, A. K. Knapp, K. L. Gross, J. E. Barrett, S. D. Frey,
638 L. Gough, R. J. Miller, J. T. Morris, L. E. Rustad, and J. Yarie. 2015. Global environmental
639 change and the nature of aboveground net primary productivity responses : insights from
640 long- term experiments. *Oecologia* 117:935–947.

641 Stevens, C. J., N. B. Dise, O. J. Mountford, and D. J. Gowing. 2004. Impact of nitrogen
642 deposition on the species richness of grasslands. *Science* 303:1876–1879.

643 Suding, K. N., S. L. Collins, L. Gough, C. Clark, E. E. Cleland, K. L. Gross, D. G. Milchunas,
644 and S. Pennings. 2005. Functional- and abundance-based mechanisms explain diversity loss

645 due to N fertilization. PNAS 102:4387–4392.

646 Tilman, D. 1982. Resource competition and community structure. Princeton University Press,
647 Princeton, NJ.

648 Tilman, D. 1987. Secondary succession and the pattern of plant dominance along experimental
649 nitrogen gradients.

650 Tilman, D., J. Knops, D. Wedin, P. Reich, M. Ritchie, and E. Siemann. 1997. The influence of
651 functional diversity and composition on ecosystem processes. Science 277:1300–1302.

652 Vicente-Serrano, S. M., S. Beguería, and J. I. López-Moreno. 2010. A multiscalar drought index
653 sensitive to global warming: The standardized precipitation evapotranspiration index.
654 Journal of Climate 23:1696–1718.

655 Villéger, S., H. Mason, and D. Mouillot. 2008. New multidimensional functional diversity
656 indices for a multifaceted framework in functional ecology. Ecology 89:2290–2301.

657 Violle, C., B. J. Enquist, B. J. McGill, L. Jiang, C. H. Albert, C. Hulshof, V. Jung, and J.
658 Messier. 2012. The return of the variance: Intraspecific variability in community ecology.
659 Trends in Ecology and Evolution 27:244–252.

660 Violle, C., W. Thuiller, N. Mouquet, F. Munoz, N. J. B. Kraft, M. W. Cadotte, S. W.
661 Livingstone, and D. Mouillot. 2017. Functional rarity: The ecology of outliers. Trends in
662 Ecology and Evolution 32:356–367.

663 Weiher, E., G. D. P. Clarke, and P. A. Keddy. 1998. Community assembly rules, morphological
664 dispersion, and the coexistence of plant species. Oikos 81:309–322.

665 Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. Plant and Soil
666 199:213–227.

667 Wilcox, K. R., Z. Shi, L. A. Gherardi, N. P. Lemoine, S. E. Koerner, D. L. Hoover, E. Bork, K.

M. Byrne, J. Cahill, S. L. Collins, S. Evans, A. K. Gilgen, P. Holub, L. Jiang, A. K. Knapp,
D. LeCain, J. Liang, P. Garcia-Palacios, J. Peñuelas, W. T. Pockman, M. D. Smith, S. Sun,
S. R. White, L. Yahdjian, K. Zhu, and Y. Luo. 2017. Asymmetric responses of primary
productivity to precipitation extremes: A synthesis of grassland precipitation manipulation
experiments. *Global Change Biology* 23:4376–4385.

Willems, J. H., R. K. Peet, and L. Bik. 1993. Changes in chalk-grassland structure and species
richness resulting from selective nutrient additions. *Journal of Vegetation Science* 4:203–
212.

Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-bares,
F. S. Chapin, J. H. C. Cornelissen, M. Diemer, J. Flexas, E. Garnier, P. K. Groom, J. Gulias,
K. Hikosaka, B. B. Lamont, T. Lee, W. Lee, C. Lusk, J. J. Midgler, M.-L. Navas, U.
Niinemets, J. Oleksyn, N. Osada, H. Poorter, P. Poot, L. Prior, V. I. Pyankov, C. Roumet, S.
C. Thomas, M. G. Tjoelker, E. J. Veneklaas, and R. Villar. 2004. The worldwide leaf
economics spectrum. *Nature* 428:821–827.

Yang, Y., D. Tilman, C. Lehman, and J. J. Trost. 2018. Sustainable intensification of high-
diversity biomass production for optimal biofuel benefits. *Nature Sustainability*.

Zhang, X., F. W. Zwiers, G. C. Hegerl, F. H. Lambert, N. P. Gillett, S. Solomon, P. A. Stott, and
T. Nozawa. 2007. Detection of human influence on twentieth-century precipitation trends.
Nature 448:461–465.

Tables and Figures

Table 1: Most metrics of community composition had continuous gradual responses to chronic resource addition. Here, we show which hypothesis (Fig. 1) each metrics followed for nitrogen fertilization (F, +14 g N m⁻² year⁻¹), irrigation (I, +25 mm H₂O week⁻¹ May-September), or the interaction of the two (F:I). Functional traits shifts were quantified both by community weighted means and non-weighted (NW) means. Metrics were considered to have rapid responses, or responses in the first year which distinguished them from ambient plots, if the standard error for the treatment at year 1 did not overlap with ambient standard errors. Rapid responses were not characterized for species-specific responses. Detailed results and estimated effects can be found in Figs. 3, 4, & S1.

Metric	H1 – Stochastic	H2 – Gradual	H3 – Abrupt	H4 – Transient	Rapid Response
<i>Community Metrics</i>					
Species Richness		F, I			F
Simpson's Diversity		F			
Evenness			F		
Functional Richness	F				F
Functional Dispersion					
Functional Distance (AW)		F, I			
Functional Distance (NW)	F				
<i>Functional Traits</i>					
SLA					F, I
Leaf N			I		I
Height		F, F:I			F, I
Proportion C3		F	I		
SLA (NW)			F, I		F, I
Leaf N (NW)				I	
Height (NW)				I	
Proportion C3 (NW)		F, I, F:I			F
<i>Functional Groups</i>					
C4 Grasses		F	I		
C3 Grasses		I	F		I
Legumes		I	F		I
Non-legume Forbs		F, I, F:I			F, I
<i>Dominant Species</i>					
<i>Andropogon gerardii</i>	F, I				

<i>Bouteloua gracilis</i>				F, I	
<i>Coreopsis palmata</i>		I			
<i>Dalea purpurea</i>		I			
<i>Liatris aspera</i>		F, I			
<i>Lupinus perennius</i>		F			
<i>Poa parentis</i>			I		
<i>Sorghastrum nutans</i>		I			
<i>Solidago rigida</i>		F, I, F:I			
<i>Schizachryum scoparium</i>	F				
Total Responses:	F:4, I:1	F:11, I:12	F:4, I:5	F: 1, I:3	F: 7, I:7

701

702

703

704

705

706

707

708




Figure 1: Graphical representation of how chronic resource addition can alter measures of community composition through time. In the simplest case where the ambient plots (black line) stay static through time, the solid red line is the hypothetical response to a chronic resource addition. Communities can either (H1) have random shifts ('stochastic effect'), (H2) increasingly deviate ('gradual change'), (H3) have a sudden, but persistent shift ('abrupt change'), or (H4) an initial effect that diminishes through time ('transient effect'). Functional richness responses are illustrated here, but we studied temporal trends in several metrics of community structure.




Figure 2: Multiple trait-based and taxonomic measures provide complementary information when characterizing community restructuring. We show several hypothetical responses of community (A), which is comprised of 9 species, to resource addition. Each circle is a species, plotted according to its trait values. Functional richness, the volume of the convex hull, is shown by the area inside of the connected species. For example, resource addition may have no effect on species richness, but decrease functional richness (B); or have no effect on functional richness, but decrease species richness (C); or decrease both species richness and functional richness (D); or decrease neither species richness nor functional richness, but shift the community to a different region of trait space, which can be captured by functional distance (E).




Figure 3: Fertilization (+14 g N/m²/year) and irrigation (+25 mm H₂O/week from May-September) tended to cause gradual and continuous effects on plant community structure compared to ambient plots. Fertilization had increasingly negative effects on both Simpson's Diversity (A) and species richness (B), but had a significant abrupt effect on evenness (C). Irrigation only had an increasingly negative effect on species richness (B). Fertilization caused a stochastic decrease in functional richness (D). Irrigation and fertilization caused treated communities to become increasingly dissimilar from ambient communities (F) and did not impact functional dispersion (E). Y-axis values are residuals from partial regressions after accounting for SPEI and biomass, error bands show ± 1 standard error, points represent observed average values.




Figure 4: Both functional group composition (A-D) and functional traits (E-H) were influenced by fertilization (+14 g N/m²/year) and irrigation (+25 mm H₂O/week from May-September). Fertilization and irrigation both had rapid impacts on functional groups(A-D), which then tended to diverge from ambient plot temporal trends. Fertilization tended to have gradual continuous effects on traits (E-H) and irrigation tended to have immediate and smaller persistent impacts. Y-axis values are residuals from partial regressions after accounting for SPEI and biomass, except for SLA, where these variables were not included in the model (Tables S4 & S5). Error bands show ± 1 standard error, and points are observed average values.




Figure 5: Abundance-weighted trait values show a clearer response to resource addition through time in multidimensional space. Upper panels show shifts in trait space (A,B), middle panels show shifts in functional group space (C,D), and lower panels show shifts in species community space (E,F). Loadings for traits (B), functional groups (D) and species (E, 10 most abundant shown) show the direction for increasing values. Axes are the top two non-metric multidimensional scaling (NMDS) axes, which represent most of the variation in trait, functional group, or taxonomic space. Traits used include specific leaf area (SLA), maximum vegetative height (Height), leaf nitrogen content per dry mass (Leaf N), and photosynthetic pathway (C3). Species abbreviations are AG: *Andropogon gerardii*, BG: *Bouteloua gracilis*, CP: *Coreopsis palmata*, DP: *Dalea purpurea*, LA: *Liatris aspera*, LP: *Lupinus perennius*, SN: *Sorghastrum nutans*, SR: *Solidago rigida*, SS: *Schizachryum scoparium*. *Poa parentis* is not shown in (F) because its arrow is in same direction as DP but much smaller in magnitude.

Appendix 1 - Supplemental Information

Table S1: Comparison of results using high N treatments (+14 g N/m²/year) only, grouped N treatments (+7 or +14 g N/m²/year), and both N treatments for species richness and functional dispersion as examples. The “*Fert*” row refers to either grouped treatments or the high N addition treatment depending on column heading. All other rows are as in Table 1 of main text. Values are effect size (standard error). *- p < 0.05, ** - p < 0.01, *** - p < 0.001

	Species Richness			Functional Dispersion		
	High N	Grouped N	Both N	High N	Grouped N	Both N
<i>Intercept</i>	22.42 (24.74) ***	20.62 (3.73) ***	20.62 (3.69) ***	0.13 (0.04) ***	0.13 (0.03)***	0.13 (0.03) ***
<i>SPEI</i>	6.78 (2.26) ***	5.69 (1.78) ***	5.69 (1.76) ***	-0.04 (0.02) ***	-0.04 (0.02)***	-0.04 (0.02) ***
<i>Biomass</i>	0.01 (0.005) ***	0.01 (0.004) ***	0.01 (0.004) ***	0.00001 (0.00004)	0.0000006 (0.00004)	0.000006 (0.00004)
<i>Year</i>	-5.07 [†] (2.28)***	-3.99 [†] (1.79)***	-3.99 [†] (1.79) ***	0.03 [†] (0.02)	0.03 [†] (0.02)**	0.03 [†] (0.02)**
<i>Fert (+7 only)</i>	---	---	-0.14 (1.39)	---	---	0.01 (0.01)
<i>Fert</i>	1.97 (1.43)	0.92(1.21)	1.97 (1.39)	-0.003 (0.001)	0.004 (0.01)	-0.003 (0.01)
<i>Irr</i>	0.94 (1.43)	0.85 (1.31)	0.85 (1.31)	0.003 (0.001)	0.007 (0.01)	0.007 (0.01)
<i>Fert(+7 only):Irr</i>	---	---	1.23 (1.61)	---	---	-0.03 (0.01) ***
<i>Fert:Irr</i>	0.40 (1.63)	0.82 (1.39)	0.40 (1.61)	-0.004 (0.01)	-0.02 (0.01) ***	-0.004 (0.01)
<i>Fert(+7 only):Year</i>	---	---	-1.74 (0.57) ***	---	---	-0.001 (0.01)
<i>Fert:Year</i>	-3.13 (0.61) ***	-2.43 (0.50) ***	-3.13 (0.57) ***	-0.003 (0.01)	-0.002 (0.01)	-0.003 (0.01)
<i>Irr:Year</i>	-1.34 (0.61) ***	-1.28 (0.47) ***	-1.28 (0.47) ***	0.003 (0.01)	0.0003 (0.004)	0.0003 (0.004)

772 *Table S2: Datasets from TRY (request IDs: 4261 & 4354) and associated references. Note that*
773 *the five datasets with the most observations are from the Midwest region or from Cedar Creek*
774 *which accounts for over 50% of the trait data used from TRY.*

Dataset	Reference	Observations
Midwestern and Southern US Herbaceous Species Trait Database	unpub.	1610
Plant traits of grassland species	(La Pierre and Smith 2015)	1538
Plant traits from Wisconsin, USA	unpub.	1050
Cedar Creek prairie plants (leaf, seed, dispersule, height, plant, root)	unpub.	671
Cedar Creek Savanna SLA, C, N Database	(Willis et al. 2010)	624
New York Old Field Plant Traits Database	(Siefert 2012)	584
The LEDA Traitbase	(Kleyer et al. 2008)	472
Plant Traits for Pinus and Juniperus Forests in Arizona	(Laughlin et al. 2010)	284
California Coastal Grassland Database	(Sandel et al. 2011)	216
The VISTA Plant Trait Database	(Garnier et al. 2007)	216
Plant Traits from LTER Matsch (Mazia), Italy	unpub.	200
FRED - Fine Root Ecology Database	(Iversen et al. 2017)	183
Eastern US Old Field Plant Traits Database	(Siefert et al. 2014)	170
Old fields of Eastern US (Siefert Data)	(Siefert et al. 2014)	170
GLOPNET - Global Plant Trait Network Database	(Wright et al. 2004)	162
Categorical Plant Traits Database	unpub.	155
Global Respiration Database	(Reich et al. 2008)	138
Rocky Mountain Biological Laboratory WSR/gradient plant traits	unpub.	119
Aboveground morphological traits of grassland species	(Abakumova et al. 2016)	110
Leaf and Whole Plant Traits Database	unpub.	97
French Alps Trait Data	(Gos et al. 2016)	89
Photosynthesis Traits Worldwide	(Maire et al. 2015)	79
Leaf Economic Traits Across Varying Environmental Conditions	(Wright and Sutton-Grier 2012)	78
French Massif Central Grassland Trait Database	(Louault et al. 2005)	72
Ecological Flora of the British Isles	(Fritter and Peat 1994)	71
Leaf and Whole Plant Traits Database	(Shipley 2002)	70
Flora Italia Functional Traits Hoard (FIFTH)	(Cerabolini et al. 2010)	69
Traits of 59 grassland species	(Schroeder-Georgi et al. 2016)	55
Grassland Plant Trait Database	(Takkis n.d.)	54
Reich-Oleksyn Global Leaf N, P Database	(Reich et al. 2009)	52
Italian Alps Plant Traits Database	(Bragazza 2009)	51
Leaf Physiology Database	(Kattge et al. 2009)	49
Global Respiration Database	(Atkin et al. 2015)	46
Leaf Allometry Dataset	(Price and Enquist 2007)	40
PLANTSdata USDA	(Green 2009)	33
Chinese Traits	(Prentice et al. 2011)	24
Plant Physiology Database	(Campbell et al. 2007)	24
Leaf Area, Dry Mass and SLA Dataset	unpub.	24
Northern mixed-grass prairie species traits - Wyoming, USA	unpub.	20
Traits of the Hungarian flora	(Lhotsky et al. 2016)	20
Plant Traits from Fynbos Forests in the Cape Region	(Onstein et al. 2014)	19
The DIRECT Plant Trait Database	(Fry et al. 2014)	19
BASECO: a floristic and ecological database of Mediterranean French flora	(Gachet et al. 2005)	18
The Netherlands Plant Height Database	unpub.	18
Leaf Structure, Venation and Economic Spectrum	(Blonder et al. 2012)	18
Photosynthesis and Leaf Characteristics Database	unpub.	18
Saskatchewan Plant Trait Database	(Guy et al. 2013)	18
Abisko & Sheffield Database	(Quested et al. 2003)	16
The Netherlands Plant Traits Database	(Ordoñez et al. 2010)	15
The Xylem/Phloem Database	(Schweingruber and Landolt 2005)	15
Sheffield Database	(Cornelissen et al. 1999)	14
Hokkaido leaf traits	(Mori et al. 2015)	13
PLANTATT - Attributes of British and Irish Plants	(Hill et al. 2004)	13
Sheffield-Iran-Spain Database	(Díaz et al. 2004)	13
The DIRECT Plant Trait Database	(Everwand et al. 2014)	11
Traits related to riparian plant invasion in South East Australia	(Catford et al. 2014)	11
Maxfield Meadow, Rocky Mountain Biological Laboratory - LMA	unpub.	10
Plant Physiology Database	(Loveys et al. 2003)	9

Plant Traits from Romania	(Ciocărlan 2000)	9
Abisko & Sheffield Database	(Cornelissen et al. 2004)	8
Leaf and Whole Plant Traits Database	(Meziane and Shipley 1999)	8
Nutrient Resorption Efficiency Database	(Vergutz et al. 2012)	8
Overton/Wright New Zealand Database	unpub.	8
TOPIC (Traits of Plants in Canada)	(Aubin et al. 2016)	7
Chinese Leaf Traits Database	(Han et al. 2004)	6
Functional Traits of Graminoids in Semi-Arid Steppes Database	(Adler et al. 2004)	6
Herbaceous Traits from the -land Island Database	(Hickler 1999)	6
Leaf N-Retention Database	(de Vries and Bardgett 2016)	6
The Americas N&P database	(Kerkhoff et al. 2006)	5
A Global Data Set of Leaf Photosynthetic Rates, Leaf N and P, and Specific Leaf Area	(Walker 2014)	4
Fonseca/Wright New South Wales Database	(Fonseca et al. 2000)	4
Leaf Traits and Seed Mass of Cover Crops	(Tribouillois et al. 2015)	4
Dispersal Traits Database	unpub.	3
Plant Coastal Dune Traits (France, Aquitaine)	unpub.	3
Functional traits explaining variation in plant life history strategies	(Adler et al. 2014)	2
Leaf and Whole Plant Traits Database	(Shipley and Lechowicz 2000)	2
Leaf Ash Content in China's Terrestrial Plants	(Han et al. 2012)	2
Leaf Traits Mount Hutt, New Zealand	(Kichenin et al. 2013)	2
Plant Physiology Database	(Atkin et al. 1997)	2
Plant Trait Database in East and South-East Asia	(Koike 2001)	2
Plant Traits of Acidic Grasslands in Central Spain	(Peco et al. 2005)	2
The Functional Ecology of Trees (FET) Database - Jena	(Wirth and Lichstein 2009)	2
Leaf and Whole Plant Traits Database	(Shipley 1995)	1
Leaf and Whole Plant Traits Database	(Shipley and Vu 2002)	1
Leaf and Whole Plant Traits Database	(Vile 2005)	1
Photosynthesis Traits Database	unpub.	1
Herbaceous Plants Traits From Southern Germany	unpub.	1
Mediterranean psammophytes	(Ciccarelli 2015)	1

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

Table S3: Percent trait coverage for height, leaf N content, and SLA. There was >89% coverage for all treatments each year. All species had a photosynthetic pathway, so coverage was 100% for all years and treatments.


Year	N	Water	Species Coverage			Biomass Coverage		
	Treatment	Treatment	Height	Leaf N	SLA	Height	Leaf N	SLA
2007	ambient	ambient	100	100	100	100	100	100
	Fertilized	ambient	99.24	99.17	100	99.94	99.99	100
	ambient	Irrigated	99.24	99.24	99.24	99.99	99.99	99.99
	Fertilized	Irrigated	100	98.26	100	100	99.93	100
2009	ambient	ambient	98.55	97.57	99.28	99.64	99.97	99.99
	Fertilized	ambient	97.63	95.20	97.63	99.93	98.89	99.93
	ambient	Irrigated	100	100	100	100	100	100
	Fertilized	Irrigated	97.88	97.96	99.07	99.95	99.99	99.86
2010	ambient	ambient	96.64	87.30	96.64	99.933	99.63	99.99
	Fertilized	ambient	98.72	90.55	98.72	98.96	99.45	99.96
	ambient	Irrigated	100	97.36	100	100	99.51	100
	Fertilized	Irrigated	98.41	89.44	98.41	99.91	98.71	99.91
2015	ambient	ambient	97.86	89.44	97.86	99.97	89.11	99.97
	Fertilized	ambient	96.01	84.91	98.03	96.52	95.19	99.97
	ambient	Irrigated	97.54	89.06	97.54	99.95	92.73	99.99
	Fertilized	Irrigated	96.75	89.66	98.72	99.82	99.03	99.99
2016	ambient	ambient	96.96	94.52	96.96	99.85	99.25	99.85
	Fertilized	ambient	94.29	90.70	94.29	98.62	97.92	98.62
	ambient	Irrigated	99.12	98.43	99.12	99.99	99.96	99.99
	Fertilized	Irrigated	98.81	94.81	98.81	99.99	99.45	99.99

Table S4: Functional group biomass responds to both irrigation and fertilization treatments through time. Columns correspond with the individual functional groups. Models were run using the proportion of biomass characterized by the specified functional groups. Rows correspond to model parameters, --- means that those parameters were not included in the model. Values are effect size (standard error) *- p< 0.05, ** - p < 0.01, *** - p < 0.001

	C4 Grass	C3 Grass	Legume	Non-legume Forb
<i>Intercept</i>	0.81 (0.11) ***	-0.09 (0.20)	0.74 (0.09)***	0.32 (.025)
<i>SPEI</i>	0.11 (0.07)	-0.16 (0.09) **	-0.122 (0.06)***	0.28 (0.18) ***
<i>Biomass</i>	0.004 (0.004)	-0.0003 (0.0002)	-0.001 (0.0004) ***	0.001 (0.0003) ***
<i>Year</i>	-0.06 (0.02) **	0.23 [†] (0.10) ***	0.04 (0.02) **	-0.11 [†] (0.12)
<i>Fert</i>	0.05 (0.06)	0.12 (0.06) ***	-0.16 (0.06) ***	0.14 (0.08) *
<i>Irr</i>	0.12 (0.06) **	0.16 (0.06) ***	-0.16 (0.04) **	0.09 (0.08)
<i>Fert:Irr</i>	-0.07 (0.07)	-0.11 (0.06) *	0.10 (0.07)	-0.15 (0.11)
<i>Fert:Year</i>	0.02 (0.01)***	0.03 (0.03)	-0.005 (0.006)	-0.18 (0.04) ***
<i>Irr:Year</i>	0.01 (0.01)	-0.06 (0.03) ***	0.02 (0.006) ***	-0.13 (0.04)***
<i>Fert:Irr:Year</i>	---	---	---	0.12 (0.06) **

Table S5: Fertilization either immediately impacts community weighted trait values or has a continuous effect whereas irrigation only has immediate and persistent effects. Output for the fixed effects of individual traits (columns), abundance weighted and not. Rows correspond to model parameters, --- means that those parameters were not included in the model. Values are effect size (standard error) *- p< 0.05, **- p < 0.01,*** - p < 0.001

	Abundance weighted				Non-Abundance weighted			
	SLA	Leaf N	Height	C3	SLA	Leaf N	Height	C3
<i>Intercept</i>	16.04 (0.44) ***	16.32 (1.06) ***	0.51 (0.09) ***	0.77 (0.11) ***	16.06 (0.33) ***	14.74 (1.49) ***	0.32 (0.05) ***	0.43 (0.10) ***
<i>SPEI</i>	----	-1.48 (0.74) **	0.10 (0.04) ***	-0.11 (0.07)	----	-1.92 (1.26)	0.13 (0.04) ***	-0.26 (0.07) ***
<i>Biomass</i>	----	-0.005 (0.004)	0.002 (0.0001) ***	-0.0004 (0.0004)	----	0.004 (0.004)	0.0006 (0.0001) ***	0.0002 (0.0004)
<i>Year</i>	-0.12 (0.04) ***	0.33 (0.24)	-0.05 [†] (0.04)	0.06 (0.02) ***	-0.12 (0.04) ***	6.61 (13.45)	-1.14 (0.45)***	0.08 (0.02) ***
<i>Year</i> ²	---	---	---	---	---	-1.07 (4.36)	0.30 (0.11) ***	---
<i>Fert</i>	1.07 (0.60) *	-0.71 (0.60)	0.06 (0.03) *	-0.06 (0.06)	1.26 (0.44) ***	-0.50 (0.42)	-0.03 (0.02)	0.06 (0.06)
<i>Irr</i>	0.92 (0.51) *	-1.80 (0.60) ***	0.04 (0.03)	-0.12 (0.06) **	0.78 (0.44) *	-1.14 (0.42) ***	0.01 (0.02)	-0.04 (0.06)
<i>Fert:Irr</i>	-0.89 (0.65)	0.31 (0.71)	-0.10 (0.04) ***	0.07 (0.07)	-0.92 (0.55)	-0.04 (0.59)	0.02 (0.02)	-0.08 (0.09)
<i>Fert:Year</i>	0.07 (0.05)	-0.02 (0.07)	-0.05 (0.02) **	-0.02 (0.007) ***	0.05 (0.05)	-2.32 (2.70)	-0.13 (0.09)	-0.04 (0.01) ***
<i>Irr:Year</i>	-0.07 (0.05)	0.07 (0.07)	-0.02 (0.02)	-0.009 (0.007)	-0.05 (0.05)	0.26 (2.70)	-0.01 (0.09)	-0.02 (0.01) ***
<i>Fert: Year</i> ²	---	---	---	---	---	0.91(2.70)	0.04 (0.09)	---
<i>Irr: Year</i> ²	---	---	---	---	---	8.32 (2.70) ***	-0.22 (0.09) ***	---
<i>Fert:Irr:Year</i>	---	---	0.07 (0.02) ***	---	---	---	---	0.03 (0.01) ***




823

Effect Size

824 *Supplemental figure 1: Fertilization continually impacts five of the six community metrics,*
825 *whereas irrigation only significantly impacts two of six. Output from mixed effects models*
826 *showing the impact of the fertilization and irrigation treatments on community metrics (panels)*
827 *through time. The y-axis correspond to model parameters: SPEI is the standardized precipitation*
828 *evapotranspiration index, Biomass is the average biomass across all plots for that year, Year*
829 *corresponds to duration of experiment (0 = 2007, 9 = 2016), † in the year row indicates that the*
830 *model was fit to the log of year, Fert is the nitrogen addition treatments and Irr is the irrigation*

treatments. The x-axis is the effect size and error bars are standard errors, or in the case of functional distance, 95% confidence intervals. *- $p < 0.05$, ** - $p < 0.01$, *** - $p < 0.001$



*Supplemental Figure 2: Fertilization causes plots to become increasingly dissimilar when considering non-weighted trait values, a stochastic effect (left). Points are means, lines are estimated from the MCMCglmm output, and error bands are the upper and lower 95% confidence intervals. The right panel shows the effect sizes and the 95% confidence interval for each parameter in the MCMCglmm model. *- $p < 0.05$, ** - $p < 0.01$, *** - $p < 0.001$*




Figure S3. Most species responded in a gradual and continuous manner to either irrigation or fertilization. Output from mixed effects models showing the impact of the fertilization and irrigation treatments on community metrics (panels) through time. The y-axis correspond to model parameters: SPEI is the standardized precipitation evapotranspiration index, Biomass is the average biomass across all plots for that year, Year corresponds to duration of experiment (0 = 2007, 9 = 2016), † in the year row indicates that the model was fit to the log of year, Fert is the nitrogen addition treatments and Irr is the irrigation treatments. The x-axis is the effect size and error bars are standard errors, or in the case of functional distance, 95% confidence intervals. * - $p < 0.05$, ** - $p < 0.01$, *** - $p < 0.001$. Species abbreviations are as follows - AG: *Andropogon gerardii*, BG: *Bouteloua gracilis*, CP: *Coreopsis palmata*, DP: *Dalea purpurea*, LA: *Liatris*

aspera, LP: *Lupinus perennius*, PP: *Poa parentis*, SN: *Sorghastrum nutans*, SR: *Solidago rigida*,
SS: *Schizachryum scoparium*.

References

- Abakumova, M., K. Zobel, A. Lepik, and M. Semchenko. 2016. Plasticity in plant functional traits is shaped by variability in neighbourhood species composition. *New Phytologist* 211:455–463.
- Adler, P. B., D. G. Milchunas, W. K. Lauenroth, O. E. Sala, and I. C. Burke. 2004. Functional traits of graminoids in semi-arid steppes: A test of grazing histories. *Journal of Applied Ecology* 41:653–663.
- Adler, P. B., R. Salguero-gómez, A. Compagnoni, J. S. Hsu, J. Ray-mukherjee, P. B. Adler, R. Salguero-gómez, A. Compagnoni, J. S. Hsu, and J. Ray-mukherjee. 2014. Functional traits explain variation in plant life history strategies. *Proceedings of the National Academy of Sciences* 111:740–745.
- Atkin, O. K., K. J. Bloomfield, P. B. Reich, M. G. Tjoelker, G. P. Asner, D. Bonal, G. Bönisch, M. Bradford, L. A. Cernusak, E. G. Cosio, D. Creek, K. Y. Crous, T. Domingues, J. S. Dukes, J. J. G. Egerton, J. R. Evans, G. D. Farquhar, N. M. Fyllas, P. P. G. Gauthier, E. Gloor, T. E. Gimeno, K. Griffin, R. Guerrieri, M. A. Heskell, C. Huntingford, F. Y. Ishida, J. Kattge, H. Lambers, M. J. Liddell, C. H. Lusk, R. E. Martin, T. C. Maximov, A. P. Maksimov, Y. Malhi, B. E. Medlyn, P. Meir, L. M. Mercado, N. Mirotchnick, D. Ng, Ü. Niinemets, O. S. O’Sullivan, O. L. Phillips, L. Poorter, P. Poot, I. C. Prentice, N. Salinas, L. M. Rowland, M. G. Ryan, S. Sitch, M. Slot, N. G. Smith, M. H. Turnbull, M. C. VanderWel, F. Valladares, E. J. Veneklaas, L. K. Weerasinghe, C. Wirth, I. J. Wright, K. Wythers, J. Xiang, S. Xiang, and J. Zaragoza-Castells. 2015. Global variability in leaf respiration among plant functional types in relation to climate and leaf traits. *New Phytologist* 206:614–636.
- Atkin, O. K., M. H. M. Westbeek, M. L. Cambridge, H. Lambers, and H. Pons. 1997. Leaf respiration in light and darkness: A comparison of slow- and fast-growing *Poa* species. *Plant Physiology* 113:961–965.
- Aubin, I., A. D. Munson, F. Cardou, P. J. Burton, N. Isabel, J. H. Pedlar, A. Paquette, A. R.

- 884 Taylor, S. Delagrangé, H. Kebli, C. Messier, B. Shipley, F. Valladares, J. Kattge, L.
885 Boisvert-Marsh, and D. McKenney. 2016. Traits to stay, traits to move: a review of
886 functional traits to assess sensitivity and adaptive capacity of temperate and boreal trees to
887 climate change. *Environmental Reviews* 24:164–186.
- 888 Blonder, B., V. Buzzard, I. Simova, L. Sloat, B. Boyle, R. Lipson, B. Aguilar-Beaucage, A.
889 Andrade, B. Barber, C. Barnes, D. Bushey, P. Cartagena, M. Chaney, K. Contreras, M. Cox,
890 M. Cueto, C. Curtis, M. Fisher, L. Furst, J. Gallegos, R. Hall, A. Hauschild, A. Jerez, N.
891 Jones, A. Klucas, A. Kono, M. Lamb, J. D. R. Matthai, C. McIntyre, J. McKenna, N.
892 Mosier, M. Navabi, A. Ochoa, L. Pace, R. Plassmann, R. Richter, B. Russakoff, H. S.
893 Aubyn, R. Stagg, M. Sterner, E. Stewart, T. T. Thompson, J. Thornton, P. J. Trujillo, T. J.
894 Volpe, and B. J. Enquist. 2012. The leaf-area shrinkage effect can bias paleoclimate and
895 ecology research. *American Journal of Botany* 99:1756–1763.
- 896 Bragazza, L. 2009. Conservation priority of Italian alpine habitats: A floristic approach based on
897 potential distribution of vascular plant species. *Biodiversity and Conservation* 18:2823–
898 2835.
- 899 Campbell, C., L. Atkinson, J. Zaragoza-Castells, M. Lundmark, O. Atkin, and V. Hurry. 2007.
900 Acclimation of photosynthesis and respiration is asynchronous in response to changes in
901 temperature regardless of plant functional group. *New Phytologist* 176:375–389.
- 902 Catford, J. A., W. K. Morris, P. A. Vesk, C. J. Gippel, and B. J. Downes. 2014. Species and
903 environmental characteristics point to flow regulation and drought as drivers of riparian
904 plant invasion. *Diversity and Distributions* 20:1084–1096.
- 905 Cerabolini, B. E. L., G. Brusa, R. M. Ceriani, R. de Andreis, A. Luzzaro, and S. Pierce. 2010.
906 Can CSR classification be generally applied outside Britain? *Plant Ecology* 210:253–261.
- 907 Ciccarelli, D. 2015. Mediterranean coastal dune vegetation: Are disturbance and stress the key
908 selective forces that drive the psammophilous succession? *Estuarine, Coastal and Shelf*
909 *Science* 165:247–253.
- 910 Ciocârlan, V. 2000. *Illustrated Flora of Romania. Pteridophyta et Spermatophyta*. Editura Ceres,
911 Bucuresti, Romania.
- 912 Cornelissen, J. H. C., N. Pérez-Harguindeguy, S. Díaz, J. P. Grime, B. Marzano, M. Cabido, F.
913 Vendramini, and B. Cerabolini. 1999. Leaf structure and defense control litter
914 decomposition rate across species and life forms in regional floras on two continents. *New*

915 Phytologist 143:191–200.

916 Cornelissen, J. H. C., H. M. Quested, D. Gwynn-Jones, R. S. P. Van Logtestijn, M. A. H. De
917 Beus, A. Kondratchuk, T. V. Callaghan, and R. Aerts. 2004. Leaf digestibility and litter
918 decomposability are related in a wide range of subarctic plant species and types. *Functional*
919 *Ecology* 18:779–786.

920 Díaz, S., J. G. Hodgson, K. Thompson, M. Cabido, J. H. C. Cornelissen, A. Jalili, J. P. Grime, F.
921 Zarrinkamar, Y. Asri, S. R. Band, S. Basconcelo, B. Hamzehee, M. Khoshnevi, A.
922 Shirvany, S. Yazdani, A. Bogaard, S. Boustani, M. Charles, M. Dehghan, V. Falczuk, A.
923 Hynd, G. Jones, E. Kowsary, S. Shaw, B. Siavash, and M. R. Zak. 2004. The plant traits
924 that drive ecosystems : Evidence from three continents. *Journal of Vegetation Science*
925 15:295–304.

926 Everwand, G., E. L. Fry, T. Eggers, and P. Manning. 2014. Seasonal variation in the capacity for
927 plant trait measures to predict grassland carbon and water fluxes. *Ecosystems* 17:1095–
928 1108.

929 Fonseca, C. R., J. M. C. Overton, B. Collins, and M. Westoby. 2000. Shifts in trait-combinations
930 along rainfall and phosphorus gradients. *Journal of Ecology* 88:964–977.

931 Fritter, A. H., and H. J. Peat. 1994. The ecological flora database. *Journal of Ecology* 82:415–
932 425.

933 Fry, E. L., S. A. Power, and P. Manning. 2014. Trait-based classification and manipulation of
934 plant functional groups for biodiversity-ecosystem function experiments. *Journal of*
935 *Vegetation Science* 25:248–261.

936 Gachet, S., E. Véla, and T. Tatoni. 2005. BASECO: A floristic and ecological database of
937 Mediterranean French flora. *Biodiversity and Conservation* 14:1023–1034.

938 Garnier, E., S. Lavorel, P. Ansquer, H. Castro, P. Cruz, J. Dolezal, O. Eriksson, C. Fortunel, H.
939 Freitas, C. Golodets, K. Grigulis, C. Jouany, E. Kazakou, J. Kigel, M. Kleyer, V. Lehsten, J.
940 Lepš, T. Meier, R. Pakeman, M. Papadimitriou, V. P. Papanastasis, H. Quested, F. Quétier,
941 M. Robson, C. Roumet, G. Rusch, C. Skarpe, M. Sternberg, J. P. Theau, A. Thébault, D.
942 Vile, and M. P. Zarovali. 2007. Assessing the effects of land-use change on plant traits,
943 communities and ecosystem functioning in grasslands: A standardized methodology and
944 lessons from an application to 11 European sites. *Annals of Botany* 99:967–985.

945 Gos, P., G. Loucougaray, M. P. Colace, C. Arnoldi, S. Gaucherand, D. Dumazel, L. Girard, S.

- Delorme, and S. Lavorel. 2016. Relative contribution of soil, management and traits to co-
variations of multiple ecosystem properties in grasslands. *Oecologia* 180:1001–1013.
- Green, W. 2009. USDA PLANTS Compilation, version 1, 09-02-02.
- Guy, A. L., J. M. Mischkolz, and E. G. Lamb. 2013. Limited effects of simulated acidic
deposition on seedling survivorship and root morphology of endemic plant taxa of the
Athabasca Sand Dunes in well-watered greenhouse trials. *Botany* 91:176–181.
- Han, W., Y. Chen, F. J. Zhao, L. Tang, R. Jiang, and F. Zhang. 2012. Floral, climatic and soil pH
controls on leaf ash content in China's terrestrial plants. *Global Ecology and Biogeography*
21:376–382.
- Han, W., J. Fang, D. Guo, and Y. Zhang. 2004. Leaf nitrogen and phosphorus stoichiometry
across 753 terrestrial plant species in China. *New Phytologist*:377–385.
- Hickler, T. 1999. Plant functional types and community characteristics along environmental
gradients on Oland's Great Alvar (Sweden). University of Lund, Sweden.
- Hill, M. O., C. D. Preston, and D. B. Roy. 2004. PLANTATT - attributes of British and Irish
Plants: status, size, life history, geography and habitat. Huntingdon: Centre for Ecology and
Hydrology.
- Iversen, C. M., M. L. McCormack, A. S. Powell, C. B. Blackwood, G. T. Freschet, J. Kattge, C.
Roumet, D. B. Stover, N. A. Soudzilovskaia, O. J. Valverde-Barrantes, P. M. van Bodegom,
and C. Violle. 2017. A global Fine-Root Ecology Database to address below-ground
challenges in plant ecology. *New Phytologist* 215:15–26.
- Kattge, J., W. Knorr, T. Raddatz, and C. Wirth. 2009. Quantifying photosynthetic capacity and
its relationship to leaf nitrogen content for global-scale terrestrial biosphere models. *Global
Change Biology* 15:976–991.
- Kerkhoff, A. J., W. F. Fagan, J. J. Elser, and B. J. Enquist. 2006. Phylogenetic and Growth Form
Variation in the Scaling of Nitrogen and Phosphorus in the Seed Plants. *The American
Naturalist* 168:E103–E122.
- Kichenin, E., D. A. Wardle, D. A. Peltzer, C. W. Morse, and T. Freschet. 2013. Contrasting
effects of plant inter- and intraspecific variation on community-level trait measures along an
environmental gradient. *Functional Ecology* 27:1254–1261.
- Kleyer, M., R. M. Bekker, I. C. Knevel, J. P. Bakker, K. Thompson, M. Sonnenschein, P.
Poschlod, J. M. Van Groenendael, L. Klimeš, J. Klimešová, S. Klotz, G. M. Rusch, M.

977 Hermy, D. Adriaens, G. Boedeltje, B. Bossuyt, A. Dannemann, P. Endels, L. Götzenberger,
 978 J. G. Hodgson, A. K. Jackel, I. Kühn, D. Kunzmann, W. A. Ozinga, C. Römermann, M.
 979 Stadler, J. Schlegelmilch, H. J. Steendam, O. Tackenberg, B. Wilmann, J. H. C.
 980 Cornelissen, O. Eriksson, E. Garnier, and B. Peco. 2008. The LEDA Traitbase: A database
 981 of life-history traits of the Northwest European flora. *Journal of Ecology* 96:1266–1274.
 982 Koike, F. 2001. Plant traits as predictors of woody species dominance in climax forest
 983 communities. *Journal of Vegetation Science* 12:327–336.
 984 Laughlin, D. C., J. J. Leppert, M. M. Moore, and C. H. Sieg. 2010. A multi-trait test of the leaf-
 985 height-seed plant strategy scheme with 133 species from a pine forest flora. *Functional*
 986 *Ecology* 24:493–501.
 987 Lhotsky, B., A. Csecserits, B. Kovács, and Z. Botta-Dukát. 2016. New plant trait records of the
 988 Hungarian flora. *Acta Botanica Hungarica* 58:397–400.
 989 Louault, F., V. D. Pillar, J. Aufrère, E. Garnier, and J.-F. Soussana. 2005. -Plant traits and
 990 functional types in response to reduced disturbance in a semi-natural grassland - Plant traits
 991 and functional types in response to reduced disturbance in a semi-natural grassland. *Journal*
 992 *of Vegetation Science* 16:151–160.
 993 Loveys, B., L. Atkinson, D. Sherlock, R. Roberts, A. Fitter, and O. Atkin. 2003. Thermal
 994 acclimation of leaf and root respiration : an investigation comparing inherently fast- and
 995 slow- growing plant species. *Global Change Biology* 9:895–910.
 996 Maire, V., I. J. Wright, I. C. Prentice, N. H. Batjes, R. Bhaskar, P. M. van Bodegom, W. K.
 997 Cornwell, D. Ellsworth, Ü. Niinemets, A. Ordóñez, P. B. Reich, and L. S. Santiago. 2015.
 998 Global effects of soil and climate on leaf photosynthetic traits and rates: Effects of soil and
 999 climate on photosynthetic traits. *Global Ecology and Biogeography*.
 1000 Meziane, D., and B. Shipley. 1999. Interacting determinants of specific leaf area in 22
 1001 herbaceous species: effects of irradiance and nutrient availability. *Plant Cell and*
 1002 *Environment* 22:447–459.
 1003 Mori, A. S., T. Shiono, T. F. Haraguchi, A. T. Ota, D. Koide, T. Ohgure, R. Kitagawa, R.
 1004 Maeshiro, T. T. Aung, T. Nakamori, Y. Hagiwara, S. Matsuoka, A. Ikeda, T. Hishi, S.
 1005 Hobara, E. Mizumachi, A. Frisch, G. Thor, S. Fujii, T. Osono, and L. Gustafsson. 2015.
 1006 Functional redundancy of multiple forest taxa along an elevational gradient: Predicting the
 1007 consequences of non-random species loss. *Journal of Biogeography* 42:1383–1396.

1008 Onstein, R. E., R. J. Carter, Y. Xing, and H. P. Linde. 2014. Diversification rate shifts in the
 1009 Cape Floristic Region: The right traits in the right place at the right time. *Perspectives in*
 1010 *plant ecology, evolution, and systematics* 16:331–340.

1011 Ordoñez, J. C., P. M. van Bodegom, J. M. Witte, R. P. Bartholomeus, J. R. van Hal, and R.
 1012 Aerts. 2010. Plant strategies in relation to resource supply in mesic to wet environments:
 1013 Does theory mirror nature? *The American Naturalist* 175:225–239.

1014 Peco, B., I. De Pablos, J. Traba, and C. Levassor. 2005. The effect of grazing abandonment on
 1015 species composition and functional traits: The case of dehesa grasslands. *Basic and Applied*
 1016 *Ecology* 6:175–183.

1017 La Pierre, K. J., and M. D. Smith. 2015. Functional trait expression of grassland species shift
 1018 with short- and long-term nutrient additions. *Plant Ecology* 216:307–318.

1019 Prentice, I. C., T. Meng, H. Wang, S. P. Harrison, J. Ni, and G. Wang. 2011. Evidence of a
 1020 universal scaling relationship for leaf CO₂ drawdown along an aridity gradient. *New*
 1021 *Phytologist* 190:169–180.

1022 Price, C. A., and B. J. Enquist. 2007. Scaling mass and morphology in leaves: an extension of the
 1023 WBE model. *Ecology* 88:1132–1141.

1024 Quested, H. M., J. H. C. Cornelissen, M. C. Press, M. C. Callaghan, R. Aerts, F. Trosien, P.
 1025 Riemann, D. Gwynn-Jones, A. Kondratchuk, and S. E. Jonasson. 2003. Decomposition of
 1026 sub-arctic plants with differing nitrogen economies: A functional role for hemiparasites.
 1027 *Ecology* 84:3209–3221.

1028 Reich, P. B., J. Oleksyn, and I. J. Wright. 2009. Leaf phosphorus influences the photosynthesis-
 1029 nitrogen relation: A cross-biome analysis of 314 species. *Oecologia* 160:207–212.

1030 Reich, P. B., M. G. Tjoelker, K. S. Pregitzer, I. J. Wright, J. Oleksyn, and J. L. Machado. 2008.
 1031 Scaling of respiration to nitrogen in leaves, stems and roots of higher land plants. *Ecology*
 1032 *Letters* 11:793–801.

1033 Sandel, B., J. D. Corbin, and M. Krupa. 2011. Using plant functional traits to guide restoration:
 1034 A case study in California coastal grassland. *Ecosphere* 2:1–16.

1035 Schroeder-Georgi, T., C. Wirth, K. Nadrowski, S. T. Meyer, L. Mommer, and A. Weigelt. 2016.
 1036 From pots to plots: Hierarchical trait-based prediction of plant performance in a mesic
 1037 grassland. *Journal of Ecology* 104:206–218.

1038 Schweingruber, F., and W. Landolt. 2005. The xylem database. wiss Federal Research Institute

WSL Updated.

Shipley, B. 1995. Structured interspecific determinants of specific leaf-area in 34 species of herbaceous angiosperms. *Functional Ecology* 9:312–319.

Shipley, B. 2002. Trade-offs between net assimilation rate and specific leaf area in determining relative growth rate: Relationship with daily irradiance. *Functional Ecology* 16:682–689.

Shipley, B., and M. J. Lechowicz. 2000. The functional co-ordination of leaf morphology, nitrogen concentration, and gas exchange in 40 wetland species. *Ecoscience* 7:183–194.

Shipley, B., and T. Vu. 2002. Dry matter content as a measure of dry matter concentration in plants and their parts. *New Phytologist* 153:359–364.

Siefert, A. 2012. Spatial patterns of functional divergence in old-field plant communities. *Oikos* 121:907–914.

Siefert, A., J. D. Fridley, and M. E. Ritchie. 2014. Community functional responses to soil and climate at multiple spatial scales: When does intraspecific variation matter? *PLoS ONE* 9.

Takkis, K. (n.d.). Changes in plant species richness and population performance in response to habitat loss and fragmentation.

Tribouillois, H., F. Fort, P. Cruz, R. Charles, O. Flores, E. Garnier, and E. Justes. 2015. A functional characterisation of a wide range of cover crop species: Growth and nitrogen acquisition rates, leaf traits and ecological strategies. *PLoS ONE* 10:1–17.

Vergutz, L., S. Manzoni, A. Porporato, R. F. Novais, and R. B. Jackson. 2012. A global database of carbon and nutrient concentrations of green and senesced leaves. Data set. Oak Ridge, Tennessee, U.S.A.

Vile, D. 2005. Significations fonctionnelle et ecologique des traits des especes vegetales: exemple dans une succession post-cultural mediterraneenne et generalisations.

de Vries, F. T., and R. D. Bardgett. 2016. Plant community controls on short-term ecosystem nitrogen retention. *New Phytologist* 210:861–874.

Walker, A. P. 2014. A global data set of leaf photosynthetic rates, leaf N and P, and specific leaf area. Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA. <http://dx.doi.org/10.3334/ORNLDAAAC/1224>.

Willis, C. G., M. Halina, C. Lehman, P. B. Reich, A. Keen, S. McCarthy, and J. Cavender-Bares. 2010. Phylogenetic community structure in Minnesota oak savanna is influenced by spatial extent and environmental variation. *Ecography* 33:565–577.

1070 Wirth, C., and J. W. Lichstein. 2009. The imprint of species turnover on old-growth forest
1071 carbon balances - Insights from a trait-based model of forest dynamics. Pages 81–113 *in* C.
1072 Wirth, G. Gleixner, and M. Heimann, editors. Old-Growth Forests: Function, Fate and
1073 Value. Springer, New York, Berlin, Heidelberg.

1074 Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-bares,
1075 F. S. Chapin, J. H. C. Cornelissen, M. Diemer, J. Flexas, E. Garnier, P. K. Groom, J. Gulias,
1076 K. Hikosaka, B. B. Lamont, T. Lee, W. Lee, C. Lusk, J. J. Midgler, M.-L. Navas, U.
1077 Niinemets, J. Oleksyn, N. Osada, H. Poorter, P. Poot, L. Prior, V. I. Pyankov, C. Roumet, S.
1078 C. Thomas, M. G. Tjoelker, E. J. Veneklaas, and R. Villar. 2004. The worldwide leaf
1079 economics spectrum. *Nature* 428:821–827.

1080 Wright, J. P., and A. Sutton-Grier. 2012. Does the leaf economic spectrum hold within local
1081 species pools across varying environmental conditions? *Functional Ecology* 26:1390–1398.
1082